


POWER CORRECTIONS AT LEP

D. WICKE

*Fachbereich Physik, Bergische Universität-GH,
Gaußstr. 20, 42097 Wuppertal, Germany
wicke@cern.ch*



The size of non-perturbative corrections to event shape observables is predicted to fall like powers of the inverse centre of mass energy. These power corrections are investigated for different observables from e^+e^- -annihilation measured at LEP as well as previous experiments. The obtained corrections are compared to other approaches and theoretical predictions. Measurements of α_s using power corrections are compared to conventional methods.

1 Introduction

The process of hadron production in e^+e^- -annihilation is usually depicted by three phases. The first part called the perturbative phase is described by perturbative QCD calculations. The second phase called fragmentation or hadronisation is usually described using Monte Carlo based models. It is widely hoped that the influence of this phase on event shape observables can be described by analytical means: Power corrections. A third phase containing hadron decays is believed to be well under control.

Power corrections arise from two different theoretical approaches: *Renormalons* and *analytical hadronisation models*. This suggests a connection between the picture of a hadronisation phase and the theoretical idea of renormalons.

The size of the correction due to the hadronisation process can be seen as a quality attribute for a specific observables. A small correction implies that this observable probes the parton structure more directly, allowing a more precise test of QCD predictions, e.g. measurements of α_s . As power corrections use few parameters for describing the hadronisation phase, it is hoped that they lead to a better understanding of its influence.

2 Simple Power Corrections

2.1 Mean Values

Means of infrared and collinear safe observables can be described by the sum of the perturbative part and the power correction term

$$\langle f \rangle = \frac{1}{\sigma_{\text{tot}}} \int f \frac{df}{d\sigma} d\sigma = \langle f_{\text{pert}} \rangle + \langle f_{\text{pow}} \rangle \quad (1)$$

where the perturbative prediction in 2nd order can be written as

$$\langle f_{\text{pert}} \rangle = A \frac{\alpha_s(\mu)}{2\pi} + \left(A \cdot 2\pi b_0 \ln \frac{\mu^2}{E_{\text{cm}}^2} + B \right) \left(\frac{\alpha_s(\mu)}{2\pi} \right)^2, \quad (2)$$

with A and B being given numbers^{1,2}, μ being the renormalisation scale and $b_0 = (33 - 2N_f)/12\pi$. To investigate the size and type of power corrections,

$$\langle f_{\text{pow}} \rangle = \frac{C_1}{E_{\text{cm}}} + \frac{C_2}{E_{\text{cm}}^2} \quad (3)$$

is used as a simple ansatz. It is useful to fix α_s in these fits to get comparable power coefficients.

The four mean event shapes in Figure 1 show qualitative agreement between the parton levels of the parton shower Monte Carlo and the second order part resulting from the fit. The corrections of all four means show $1/E_{\text{cm}}$ behaviour with only $1 - T$ having large χ^2 due to inconsistent data. Although the C_2 coefficients of $\langle 1 - T \rangle$ and $\langle B_{\text{sum}} \rangle$ are not consistent with 0, the dominant contributions comes from the $1/E_{\text{cm}}$ term. $\langle 1 - T \rangle$ and $\langle B_{\text{sum}} \rangle$ have larger corrections than $\langle M_{\text{h}}^2/E_{\text{vis}}^2 \rangle$ and $\langle B_{\text{max}} \rangle$. The numbers are given in Table 1.

2.2 Cut Integrals and Higher Moments

To investigate whether the non-perturbative correction of an event depends on specific values of an observable or not cut integrals $\langle x \rangle_{a\dots b} := \frac{1}{\sigma_{\text{tot}}} \int_a^b \frac{d\sigma}{dx} dx$ were investigated. While $\langle 1 - T \rangle_{0.1\dots 0.5}$ shows dominantly $1/E_{\text{cm}}$ correction like the full mean, $\langle M_{\text{h}}^2/E_{\text{vis}}^2 \rangle_{0.1\dots 0.5}$ acquires a $1/E_{\text{cm}}^2$ behaviour. (Table 1 and Figure 3a).

Another way of emphasising different ranges is to investigate higher moments. The OPAL Coll. investigated the power dependence of the first three moments of $1 - T$ and the C -parameter⁷.

$$\langle f_{\text{pow}} \rangle = a_f \frac{1}{n^2} \left(\frac{\mu_I}{E_{\text{cm}}} \right)^n \left[\bar{\alpha}_{n-1}(\mu_I) - \alpha_s(\mu) - \left(b_0 \cdot \ln \frac{\mu^2}{\mu_I^2} + \frac{K}{2\pi} + \frac{2b_0}{n} \right) \alpha_s^2(\mu) \right], \quad (4)$$

was used as power term. $\bar{\alpha}_{n-1}$ being a non-perturbative parameter accounting for the contributions to the event shape below an infrared matching scale μ_I , $K = (67/18 - \pi^2/6)C_A - 5N_f/9$ and $a_f = 4C_f/\pi$. Beside α_s these formulae contain $\bar{\alpha}_{n-1}$ as the only free parameters. The results shown for $\langle 1 - T \rangle$ in Figure 2 and Table 2 indicate that the assumed power law of $1/E_{\text{cm}}^n$ for the n -th moment does work.

Table 1: Coefficients from fitting Equations (1–3) to various observables with $\alpha_s(M_Z) = 0.118$ fixed.

Observable	C_1 [GeV]	C_2 [GeV ²]	$\frac{\chi^2}{\text{ndf}}$	Observable	C_1 [GeV]	C_2 [GeV ²]	$\frac{\chi^2}{\text{ndf}}$
$\langle 1 - T \rangle$	1.09 ± 0.03	-3.2 ± 0.9	72/38	$\langle 1 - T \rangle_{0.2\dots 0.5}$	0.37 ± 0.03	0.6 ± 1.2	20/9
$\langle M_{\text{h}}^2/E_{\text{vis}}^2 \rangle$	0.67 ± 0.12	-1.9 ± 0.7	21/25				
$\langle B_{\text{sum}} \rangle$	1.27 ± 0.09	-10.3 ± 4.4	21/19	$\langle \frac{M_{\text{h}}^2}{E_{\text{vis}}^2} \rangle_{0.1\dots 0.5}$	0.08 ± 0.03	9.2 ± 0.9	8/5
$\langle B_{\text{max}} \rangle$	0.33 ± 0.06	0.2 ± 2.3	23/20				




Figure 1: Measured $\langle 1 - T \rangle$, $\langle M_h^2/E_{vis}^2 \rangle$, $\langle B_{sum} \rangle$ and $\langle B_{max} \rangle^{3,4,5,6}$ as a function of the centre of mass energy. On the lefthand-side the solid lines present the hadron level prediction of JETSET74, the dotted lines show the parton shower result. On the righthand-side the solid lines present the results of the fits with Eqs. (1-3), the dotted lines show the perturbative part only. The strong decrease of the $\langle B_{sum} \rangle$ prediction at low energies at the right is caused by the large and negative C_2 coefficient which is poorly determined.




Figure 2: Results of fitting Eqs. (1,2,4) to JETSET predictions for mean values and higher moments of $1 - T$.

Table 2: Results of fitting Eqs. (1,2,4) to JETSET predictions for mean values and higher moments of $1 - T$.

$1 - T$	Mean	2nd Moment	3rd Moment
$\alpha_s(M_Z)$	0.108	0.1121	0.1131
$\bar{\alpha}_{n-1}$	0.380	1.34	5.7
χ^2/ndf	16	4.8	3.0

3 Other Approaches

To strengthen the need for power corrections in the description of data, a few other approaches were investigated: Fitting only the perturbative prediction including a 3rd order coefficient as free parameter yields very large χ^2 and a 3rd order coefficient $C = \mathcal{O}(1000)$ for $\langle 1 - T \rangle$ and $\langle M_h^2/E_{\text{vis}}^2 \rangle$. Thus a 3rd order calculation does not give an improved prediction on the energy dependence neglecting hadronisation effects. (Figure 3b).

Leaving the renormalisation scale $x_\mu = \mu^2/E_{\text{cm}}^2$ free, shows ambivalent results. While for $\langle 1 - T \rangle$ the fit is reasonable and leads to a scale x_μ that is consistent with the one obtained from fits to distributions, the fit for $\langle M_h^2/E_{\text{vis}}^2 \rangle$ cannot describe the data. (Figure 3c).

To obtain the functional type of power corrections needed, one can also fit with an arbitrary power p :

$$\langle f_{\text{pow}} \rangle = \frac{C_p}{E_{\text{cm}}^p} \quad (5)$$

For $\langle 1 - T \rangle$ one gets $p = 0.98 \pm 0.19^8$ in perfect agreement with the previous results.

4 Determination of α_s using Power Corrections

The analytical power ansatz for non-perturbative corrections by Dokshitzer and Webber^{9,10} is used by DELPHI to determine α_s from mean event shapes^{5,6}. The power correction of this ansatz is given by

$$\langle f_{\text{pow}} \rangle = a_f \cdot \frac{\mu_I}{E_{\text{cm}}} \left[\bar{\alpha}_0(\mu_I) - \alpha_s(\mu) - \left(b_0 \cdot \ln \frac{\mu^2}{\mu_I^2} + \frac{K}{2\pi} + 2b_0 \right) \alpha_s^2(\mu) \right] \quad , \quad (6)$$

with μ_I , K , a_f and b_0 as given above. In order to measure α_s from individual high energy data the free parameter $\bar{\alpha}_0$ has to be known.

To infer $\bar{\alpha}_0$ a combined fit of α_s and $\bar{\alpha}_0$ to a large set of measurements at different energies³ is performed. For $E_{\text{cm}} \geq M_Z$ only DELPHI measurements are included in the fit. The resulting values of $\bar{\alpha}_0$ are summarised in Table 3. The extracted $\bar{\alpha}_0$ values are around 0.5 as expected in¹⁰. The numerical values are, however, incompatible with each other. So the assumed universality¹⁰ is not valid to the precision that is accessible from the data. Therefore $\bar{\alpha}_0$ is determined for $\langle 1 - T \rangle$ and $\langle M_h^2/E_{\text{vis}}^2 \rangle$ individually. The scale error is obtained from varying the renormalisation scale factor $x_\mu = \mu^2/E_{\text{cm}}^2$ from 0.25 to 4 and the infrared matching scale from 1 GeV to 3 GeV.

After having fixed $\bar{\alpha}_0$, the α_s values corresponding to the high energy data points can be calculated from Eqs. (1,2,6). α_s is calculated for both observables individually and then combined with an unweighted average. The resulting α_s values and the QCD expectation are shown in the leftmost plot of Figure 4.

The α_s values follow the QCD expectation. Fitting a straight line to the energy dependence results in a slope, which agrees very well with the QCD expectation of a running α_s between 91 GeV and 183 GeV. In Figure 4 and Table 4 this result is further-on compared to α_s measurements obtained from distributions using Monte Carlo based hadronisation models. The comparison shows that the running measured from means using power correction and the one obtained from standard methods give consistent results and comparable errors.

5 Prospects

Many of the recent developments in the field of power corrections are not yet included in the presented experimental works. Beside the calculation of Milan factors, which will not influence the consistency of the experimental results much, some of the predicted power coefficients a_f were corrected in the last months¹¹. First tests show that the consistency of measured $\bar{\alpha}_0$ and α_s values improve with these new predictions.

Thus the experimental fits have to be repeated and with increasing theoretical understanding consistent experimental results hopefully will arise.

References

1. R. K. Ellis, D. A. Ross, and A. E. Terrano. *Nucl. Phys* **B178**(1981) 412.
2. CERN 89-08 Vol. 1, 1989.
3. ALEPH Coll. *Phys. Lett.* **B284** (1992) 163; ALEPH Coll. *Z. Phys.* **C55** (1992) 209; AMY Coll. *Phys. Rev. Lett.* **62** (1989) 1713; AMY Coll. *Phys. Rev.* **D41** (1990) 2675; CELLO Coll. *Z. Phys.* **C44** (1989) 63; HRS Coll. *Phys. Rev.* **D31** (1985) 1; JADE Coll. *Z. Phys.* **C25** (1984) 231; JADE Coll. *Z. Phys.* **C33** (1986) 23; P.A. Movilla Fernandez, et. al. and the JADE Coll. *Eur. Phys. J.* **C1** (1998) 461; L3 Coll. *Z. Phys.* **C55** (1992) 39; Mark II Coll. *Phys. Rev.* **D37** (1988) 1; Mark II Coll. *Z. Phys.* **C43** (1989) 325; MARK J Coll. *Phys. Rev. Lett.* **43** (1979) 831; OPAL Coll. *Z. Phys.* **C59** (1993) 1; PLUTO Coll. *Z. Phys.* **C12** (1982) 297; SLD Coll. *Phys. Rev.* **D51** (1995) 962; TASSO Coll. *Phys. Lett.* **B214** (1988) 293; TASSO Coll. *Z. Phys.* **C45** (1989) 11; TASSO Coll. *Z. Phys.* **C47** (1990) 187; TOPAZ Coll. *Phys. Lett.* **B227** (1989) 495; TOPAZ Coll. *Phys. Lett.* **B278** (1992) 506; TOPAZ Coll. *Phys. Lett.* **B313** (1993) 475.
4. L3 Coll. M. Acciarri et al. *Phys. Lett.* **B411**(1997) 339.
5. J. Drees, A. Grefrath, K. Hamacher, O. Passon, and D. Wicke. DELPHI 97-92 CONF 77, contributed to the EPS HEP conference in Jerusalem, 1997.
6. J. Drees, U. Flagmeyer, K. Hamacher, O. Passon, R. Reinhardt, and D. Wicke. DELPHI 98-18 CONF 119, contributed to Moriond, 1998.
7. OPAL Coll. OPAL PN 310, 1997.
8. ALEPH Coll. LP 258, Contributed to Lepton-Photon Conference, Hamburg, 1997.
9. Y. L. Dokshitzer and B. R. Webber. *Phys. Lett.* **B352**(1995) 451.
10. B. R. Webber. Talk given at workshop on DIS and QCD in Paris, hep-ph/9510283, 1995.
11. A. Lucenti and G. Salam. In these proceedings.




Figure 3: Fit results for cut variables (left), fitted 3rd order coefficient (middle) and fitted renormalisation scale (right). The continuous line represents the complete prediction. In the left plot the dashed line gives the perturbative contribution. In the middle plot it represents the 2nd order contribution to the full 3rd order curve.

Table 3: Determination of $\bar{\alpha}_0$. For $E_{cm} \geq M_Z$ only DELPHI measurements are included in the fit. The first error is the statistical error from the fit, the second one is the scale error.

Observable	$\bar{\alpha}_0$	$\alpha_s(M_Z)$	$\Lambda_{\overline{MS}}[\text{MeV}]$	χ^2/ndf
$\langle 1 - T \rangle$	$0.531 \pm 0.012 \pm 0.003$	$0.1187 \pm 0.0017 \pm 0.0060$	$235 \pm 22 \pm 70$	42/23
$\langle M_h^2/E_{vis}^2 \rangle$	$0.434 \pm 0.010 \pm 0.010$	$0.1144 \pm 0.0012 \pm 0.0041$	$184 \pm 13 \pm 40$	4.0/14




Figure 4: Energy dependence of α_s as obtained from mean event shapes (left) compared to α_s obtained from distributions. The errors shown are statistical only. The band shows the QCD expectation of extrapolating the world average to other energies.

Table 4: Running of α_s ($d\alpha/dE_{cm}$) between 91 GeV and 183 GeV.

Means plus Power corr. $\mathcal{O}(\alpha_s^2)$	Distributions plus Monte Carlo based hadronisation models			QCD pred.
	$\mathcal{O}(\alpha_s^2)$	NLLA	$\mathcal{O}(\alpha_s^2)+\text{NLLA}$	
-0.13 ± 0.05	-0.13 ± 0.05	-0.15 ± 0.05	-0.13 ± 0.04	$-0.13 \cdot 10^{-3} \text{ GeV}^{-1}$